

UNCLASSIFIED

AD NUMBER
AD843596
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Foreign Government Information; AUG 1968. Other requests shall be referred to the Army Biological Laboratory, Attn: Technical Release Branch [TID], Fort Detrick, MD 21701.
AUTHORITY
bdrl, per d/a ltr dtd 29 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD 843596

AD

TECHNICAL MANUSCRIPT 473

A COMPARISON OF THE IN VITRO VIABILITY
OF OVA FROM HORMONE-TREATED MICE
WITH OVA FROM SPONTANEOUS OVULATIONS

James R. Spears

Jerry S. Walker

AUGUST 1968

STATEMENT OF CONFIDENTIALITY

This document is a type of technical report
control, and its contents may be made
available to foreign nationals only with the
approval of the Department of the Army,
Fort Detrick, AFAT (Technical Release Branch,
TID, Frederick, Maryland 21721).

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

NOV 26 1968

PERMISSION FOR	
CPSTI	WHITE SECTION <input type="checkbox"/>
DOC	DIFF SECTION <input checked="" type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
DISSEMINATION	
Y	
DISTRIBUTION AVAILABILITY CODES	
8467	AVAIL. CODE SPECIAL
2	

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 473

A COMPARISON OF THE IN VITRO VIABILITY OF OVA FROM HORMONE-TREATED MICE
WITH OVA FROM SPONTANEOUS OVULATIONS

James R. Spears

Jerry S. Walker

Process Development Division
AGENT DEVELOPMENT AND ENGINEERING LABORATORY

Project 1B533001D426

August 1968

In conducting the research described in this report, the investigators adhered to the "Guide for Laboratory Animal Facilities and Care," as promulgated by the Committee on the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Academy of Sciences-National Research Council.

ACKNOWLEDGMENT

The authors acknowledge the constructive comments and suggestions of Dr. Ralph E. Lincoln in the preparation of the manuscript and Mrs. Shirley Snowden's assistance with the statistical analysis of the data.

ABSTRACT

The results of this study show that the viability of ova obtained from hormone-treated females is equal to that of ova from spontaneous ovulations in the two strains of mice studied (Swiss-Webster and C57BAL/6Cum). Approximately 90% of all cultured ova developed from the late two-cell or early four-cell stages to blastocysts. The use of exogenous hormones to induce ovulation seems to afford a reliable, efficient, and economic means of obtaining fertilized mammalian ova.

I. INTRODUCTION

The technique of inducing superovulation has afforded increased opportunity for studies using preimplantation mammalian ova. This technique provides a convenient supply of ova for ovum transfer and culture experiments. Not only are larger numbers of ova recovered per female, but, more importantly, the stages of maturation division, ovulation, and the first and subsequent cleavage divisions can be ascertained following treatment with gonadotropins.¹

Gates² concluded that the developmental capacity of eggs from immature mice treated with gonadotropins closely approximated that of eggs from spontaneous ovulations in adult mice. In his experiment, eggs from immature donors were compared with spontaneously ovulated, untransplanted eggs developing in the same uterine environment.

Chang³ obtained 53 two-cell ova from a single rabbit doe after treatment with gonadotropins, and they were transplanted into four recipient females and yielded 45 normal young. Dowling⁴ likewise reported a high incidence of success utilizing ova from superovulated does for ovum transfer. Willett, Buckner, and Larson⁵ reported three successful transplantations of fertilized bovine eggs that were obtained from superovulated females.

Brinster⁶⁻¹⁰ utilized the superovulation technique in mice to obtain a very large number of two-cell ova that developed to blastocysts with a high incidence of success. Data accumulated from studies utilizing ova from superovulated animals for ovum transfer culture^{4,5,10,11} and metabolic studies^{6-10,12,13} indicate their viability is equal to that of ova from spontaneous ovulations. However, specific studies comparing the in vitro development of spontaneous and superovulated ova have not been conducted.

This paper compares the viability of ova from hormone-induced ovulations with that of ova from spontaneous ovulations. Late two-cell and early four-cell ova from two strains of mice were tested for their ability to develop to blastocysts in vitro.

II. MATERIALS AND METHODS

A. ANIMALS

The Fort Detrick strain of Swiss-Webster mice obtained from the Fort Detrick animal farm and C57BAL/6Cum mice obtained from Cumberland View Farms, Clinton, Tennessee, were utilized in this study. All animals were 6 to 8 weeks of age at the time of use.

B. OVULATION INDUCTION

Ova were recovered from hormone-treated females induced to ovulate by an intraperitoneal injection of 5 international units (IU) of pregnant mares' serum (Ayerst) followed 44 to 50 hours later with 5 IU of human chorionic gonadotropin (HCG) (Organon). Females were placed with males immediately after the time of luteinizing hormone injection. Females exhibiting vaginal plugs (approximately 90%) the following morning were considered to be pregnant. Ova were recovered 48 to 52 hours after HCG injection.

The ova from spontaneous ovulations were recovered late on day 2 of pregnancy (day of vaginal plug is considered to be day 1 of pregnancy). This was 40 to 44 hours after the presumed time of ovulation.

C. OVA RECOVERY

The animals were sacrificed by cervical separation. The oviducts were cut free of the ovary and uterus and placed in 0.5 to 1.0 ml of culture medium (see below) in a capped plastic test tube (17 by 60 mm). The ova were recovered by inserting a blunted 30-gauge needle attached to a 2-ml syringe into the fimbrial end of the oviduct and flushing the contents into a small watch glass. The ova were manipulated by means of a pipette constructed from a drawn-out capillary tube inserted into a 16-gauge needle attached to a 1-ml Hamilton thumb-wheel syringe. All manipulations were carried out with the aid of a stereomicroscope at 30 to 60X magnification.

D. MEDIUM PREPARATION

The medium and culture techniques are modifications of those presented by Brinster.⁶ The components and concentrations of the culture medium are given in Table 1. The medium was prepared in either 100- or 200-ml quantities and stored at 4 C until ready for use. To each 100 ml of 1X Earle's balanced salt solution (EBSS), commercial preparation without bicarbonate, was added 0.2 ml of 60% sodium lactate and 1.0 ml of a 0.3% solution of

sodium pyruvate. Penicillin, streptomycin, and bovine plasma albumin were also added before storage. At the time of use, 10 ml of the above stock solution was placed in a 30-ml T-30 Falcon tissue culture flask and sodium bicarbonate was added (1.3 ml of a 1.3% solution). This 10-ml aliquot was sparged for 10 minutes with 5% CO₂ in air.

TABLE 1. MEDIUM FOR IN VITRO CULTIVATION OF MOUSE OVA

Component	Grams/Liter	
NaCl	6.8	} Earle's Balanced Salt Solution
KCl	0.4	
NaH ₂ PO ₄	0.125	
MgSO ₄ ·7H ₂ O	0.2	
CaCl ₂	0.2	
Glucose	1.0	
Phenol Red	0.02	
Sodium lactate	1.412	
Sodium pyruvate	0.03	
Bovine plasma albumin	1.00	
Penicillin G (potassium)	100 U/ml	
Streptomycin sulfate	50 µg/ml	
NaHCO ₃ ^{a/}	1.7	

a. Sodium bicarbonate is added as a 1.3% solution.

E. OVA CULTURE

Ova were individually handled and deposited in microdrops (0.05 to 0.1 ml) of culture medium in the center well of an organ culture dish (Falcon, 15 by 60 mm) under light weight paraffin oil (Saybolt viscosity 125/135, Fisher). The paraffin oil was equilibrated with the medium by adding 10 ml of the medium to 300 ml of the sterile paraffin oil and sparging for 15 minutes with 5% CO₂ in air. The mixture was stored at least 48 hours in a CO₂ incubator (Hotpack, #528) (Fig. 1) before use. A Plexiglas compartment was constructed and installed in the incubator (Fig. 2). This facilitated the retention and reequilibration of the CO₂ atmosphere upon opening the door to the incubator. The gas flow in the incubator was controlled by standard flow meters on the incubator.




Figure 1. Hotpack Incubator #528.

The culture dishes containing the ova in the microdrops were placed in the incubator within 8 to 10 minutes after the mice were sacrificed. Whenever desired, the culture dishes were removed and observed or photographed.

For an ovum to be selected or distinguished as normal, it had to meet the following criteria: fertilized, zona pellucida intact, blastomeres of approximately equal size, and no other gross morphological abnormality. Ova not meeting one or more of these criteria were considered abnormal and were discarded immediately after recovery.



Figure 2. Plexiglas Compartment Installed in the Incubator to Facilitate Retention and Reequilibration of CO₂ Atmosphere.

The number of normal ova developing to blastocyst after 3 days in culture was determined microscopically using an inverted microscope (1 to 200X). The proportions of those ova developing to blastocyst were determined from the total number cultured at any one time.

All cultures in which less than 50% of the ova developed to blastocysts were excluded because percentages this low are indicative of developmental failures attributable to factors other than those being studied.

III. RESULTS

A total of 879 mouse ova were cultured in vitro, and 88% developed from the late two-cell or early four-cell stages (Fig. 3) to blastocyst (Fig. 4). These results are in accord with those presented by Brinster,⁸ who demonstrated that 60 to 100% of the two-cell ova recovered 45 hours after HCG injection developed to blastocyst.

In the present study, the ability of ova from hormone-treated females of two strains of mice to develop from early preimplantation stages to blastocysts in vitro was compared with that of ova of equal development from spontaneous ovulations. The numbers and proportions of ova developing to blastocysts are presented in Table 2. Preliminary investigations indicated that the developmental capacity of late two-cell ova did not differ appreciably from that of early four-cell ova. In view of these preliminary findings, no distinction was made between the number of four-cell ova recovered (25% of the total) and two-cell ova in Table 2. The data were transformed by determining the square roots of the percentages for normalizing purposes and analyzed by means of the "t" test to compare sample means. There was no significant difference between any of the groups tested at $P \leq 0.05$ level.




Figure 3. Two- and Four-Cell Mouse Ova at Beginning of In Vitro Culture. 200X.



Figure 4. Blastocysts Developed from Two- and Four-Cell Ova Cultured In Vitro for 48 Hours. 200X.

TABLE 2. PROPORTIONS OF OVA DEVELOPING FROM TWO- AND FOUR-CELL STAGE TO BLASTOCYST RECOVERED
FROM HORMONE-TREATED AND SPONTANEOUSLY OVULATING SWISS-WEBSTER
AND C₅₇BAL/6CUM FEMALE MICE

Hormone-Induced Ovulation				Spontaneous Ovulation			
C ₅₇		Swiss-Webster		C ₅₇		Swiss-Webster	
No. Developed/ Total No.	%	No. Developed/ Total No.	%	No. Developed/ Total No.	%	No. Developed/ Total No.	%
8/15	53.3	7/9	77.8	3/3	100.0	3/3	100.0
11/13	84.6	7/8	87.5	3/3	100.0	10/10	100.0
25/31	80.6	4/4	100.0	9/11	81.8	9/9	100.0
5/7	70.4	14/15	93.3	7/8	87.5	10/11	90.9
3/3	100.0	13/15	86.7	9/9	100.0	8/10	80.0
28/28	100.0	10/16	60.0	6/6	100.0	5/6	83.3
6/7	85.7	7/10	70.0	6/7	85.7	4/6	66.7
35/37	94.7	8/10	80.0	8/9	88.9	15/15	100.0
30/35	85.7	21/23	91.3	13/14	92.8	10/10	100.0
14/15	93.3	19/22	86.4	14/14	100.0	8/9	88.9
22/24	91.7	12/13	92.3	5/7	71.4	11/14	78.9
4/7	57.1	29/32	90.6	8/9	88.9	7/9	77.8
16/16	100.0	20/23	86.9	8/8	100.0	9/10	90.0
4/4	100.0	18/24	75.0	-	-	28/28	100.0
6/6	100.0	14/16	87.5	-	-	-	-
4/4	100.0	14/16	87.5	-	-	-	-
5/7	71.4	15/16	93.8	-	-	-	-
2/2	100.0	16/18	88.9	-	-	-	-
5/5	100.0	-	-	-	-	-	-
25/27	92.6	-	-	-	-	-	-
10/10	100.0	-	-	-	-	-	-
24/26	92.3	-	-	-	-	-	-
Total							
292/331	88.2	248/290	85.5	99/108	91.7	137/150	91.3
Overall Total							
776/879 = 88%							

IV. DISCUSSION

Admittedly, criticism can be directed toward the use of both two- and four-cell ova in this experiment. However, it seemed of utmost importance to ensure that those ova from spontaneous ovulations developed sufficiently in vivo to allow continued development in vitro. This meant that ova recovered from spontaneous ovulations had to remain in the oviduct at least 32 to 35 hours after ovulation. The presence of four-cell ova some 40 to 44 hours after the presumed time of ovulation was a true indication that sufficient time had elapsed. In the initial stages of this experiment, the four-cell ova were cultured separately and the proportions recorded. The developmental efficiency of these ova did not exceed that of two-cell ova recovered at the same time. Thus, no distinction is made between two- and four-cell ova in Table 2.

The recovery of ova from hormone-treated females was performed at a sufficient time interval after HCG injection to ensure that some of these ova also reached the four-cell stage. As noted in Section II, recoveries performed 48 to 52 hours after HCG injection exhibited a preponderance of two-cell ova.

The culture technique, as mentioned previously, is a modification of that described by Brinster;⁶ it is important to point out the simplicity of his technique. By using commercially prepared salts (EBSS), the medium can be prepared in a very few minutes and in small quantities. This medium contains not only lactate and pyruvate as energy sources, but also glucose, which is in the EBSS. Isotonic sodium bicarbonate solution (1.3%) was utilized for pH adjustment. Preliminary studies showed that 1.3 ml is the optimum quantity of bicarbonate solution to add to the 10 ml of the medium. This produces a pH of approximately 6.8 after equilibration.

Studies involving the use of preimplantation mammalian ova offer great potential for cell research. Investigations conducted with such cells could provide insight into factors influencing differentiation, metabolic pathways, blastomere potency, and the effects of viruses upon the earliest embryonic development. The establishment of permanent cell lines from these cells would most certainly provide a useful research tool.

LITERATURE CITED

1. Edwards, R.G.; Gates, A.H. 1959. Timing of the stages of the maturation divisions, ovulation, fertilization and the first cleavage of eggs of adult mice treated with gonadotropins. J. Endocrinol. 18:292-304.
2. Gates, A.H. 1956. Viability and developmental capacity of eggs from immature mice treated with gonadotropins. Nature 177:754-755.
3. Chang, M.C. 1948. Probability of normal development after transplantation of fertilized rabbit ova stored at different temperatures. Proc. Soc. Exp. Biol. Med. 68:680-683.
4. Dowling, D.F. 1949. Problem of the transplantation of fertilized ova. J. Agr. Sci. 39:374-396.
5. Willett, E.L.; Buckner, P.J.; Larson, G.L. 1953. Three successful transplantations of fertilized bovine eggs. J. Dairy Sci. 36:520-523.
6. Brinster, R.L. 1963. A method for in vitro cultivation of mouse ova from two-cell to blastocyst. Exp. Cell Res. 32:205-208.
7. Brinster, R.L. 1965. Studies on the development of mouse embryos in vitro: I. The effect of osmolarity and hydrogen ion concentration. J. Exp. Zool. 158:49-58.
8. Brinster, R.L. 1965. Studies on the development of mouse embryos in vitro: II. The effect of energy source. J. Exp. Zool. 158:59-68.
9. Brinster, R.L. 1965. Studies on the development of mouse embryos in vitro: III. The effect of fixed nitrogen source. J. Exp. Zool. 158:69-78.
10. Brinster, R.L. 1966. Malic dehydrogenase activity in the preimplantation mouse embryo. Exp. Cell Res. 43:131-135.
11. Willett, E.L.; Black, W.G.; Casida, L.E.; Stone, W.H.; Buckner, P.J. 1951. Successful transplantation of a fertilized bovine ovum. Science 113:247.
12. Auerbach, S.L.; Brinster, R.L. 1967. Lactate dehydrogenase isozymes in the early mouse embryo. Exp. Cell Res. 46:89-92.
13. Thompson, J.L. 1967. Effect of inhibitors of carbohydrate metabolism on the development of preimplantation mouse embryos. Exp. Cell Res. 46:252-262.

Unclassified
Security Classification

15

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
Department of the Army Fort Detrick, Frederick, Maryland 21701		Unclassified
		2b. GROUP
3. REPORT TITLE		
A COMPARISON OF THE IN VITRO VIABILITY OF OVA FROM HORMONE-TREATED MICE WITH OVA FROM SPONTANEOUS OVULATIONS		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name)		
James R. Spears Jerry S. Walker		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
August 1968	15	13
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S)
A. PROJECT NO. 1B533001D426		Technical Manuscript 473
9.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
10. DISTRIBUTION STATEMENT		
Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY
		Department of the Army Fort Detrick, Frederick, Maryland 21701
13. ABSTRACT		
The results of this study show that the viability of ova obtained from hormone-treated females is equal to that of ova from spontaneous ovulations in the two strains of mice studied (Swiss-Webster and C57BAL/6Cum). Approximately 90% of all cultured ova developed from the late two-cell or early four-cell stages to blastocysts. The use of exogenous hormones to induce ovulation seems to afford a reliable, efficient, and economic means of obtaining fertilized mammalian ova.		
14. Key Words		
Ova Superovulation Gonadotropins *Viability Mice		

DD FORM 1473 REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified
Security Classification